


AD 697 864

# BIOLOGICAL CHARACTERISTICS OF UNA VIRUS

O. N. Shcheglovitova


Army Foreign Science and Technology Center  
Washington, D. C.

27 October 1969



This document has been approved for public release and sale.

# DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

AD 697 864

FSTC-HT-23-061-70

# U.S. ARMY FOREIGN SCIENCE AND TECHNOLOGY CENTER




BIOLOGICAL CHARACTERISTICS OF UNA VIRUS

COUNTRY: USSR

## TECHNICAL TRANSLATION

Reproduced by the  
CLEARINGHOUSE  
for Federal Scientific & Technical  
Information Springfield Va. 22151

This document has been approved  
for public release and sale;  
its distribution is unlimited.  
It may be released to the  
Clearinghouse; Department of  
Commerce, for sale to the  
general public.



INDEXED ☐ ☐  
 WRITE SECTION ☐  
 COPY SECTION ☐  
 MAGNIFIED ☐  
 DATE-18-1988  
 BY \_\_\_\_\_  
 DISTRIBUTION/AVAILABILITY CODES  
 GUCI \_\_\_\_\_  
 7

**TECHNICAL TRANSLATION**

**FSTC-HT-23- 061-70**

**BIOLOGICAL CHARACTERISTICS OF UNA VIRUS**

by

**O. N. Shcheglovitova**

**Source:**  
**Ivanovskiy Institute of Virology Publication**  
**Pp. 221-224**  
**USSR**

**Translated ofr FSTC by ACSI**

This translation is an unedited rendition of the original foreign text. Statements or theories advocated or implied are those of the source and do not reflect the position or opinion of the US Army Foreign Science and Technology Center. This translation is published with a minimum of copy editing and graphics preparation in order to expedite the dissemination of information. Requests for additional copies of this document should be addressed to the Defense Documentation Center, Cameron Station, Alexandria, Virginia, ATTN: DoR-1.

**BIOLOGICAL CHARACTERISTICS OF UNA VIRUS**  
(Report 2)

Ivanovskiy Institute of Virology  
Publication  
Pages 221-224

O.N. Shcheglovitova

Interest in arbor viruses has heightened recently in view of the isolation of such viruses in many of the world's countries. The present investigation deals with the biological distinctions and development of optimum methods for passing and preserving Una arbor virus which is referable to group A according to the Casals classification. In the course of this study it was found that Una virus induces CPE [cytopathogenic effect] in VNK-21 tissue cultures and plaque production in chick fibroblast cultures. The clinical symptoms of pathology following infection of albino mice weighing 6-7 grams with Una virus, intracerebrally, are characterized by acute development of disease, paralysis of the hind legs on the 4th-5th postinfection day. The virus is demonstrable in the blood in the highest titers 6 and 48 hours after infection, and in the brain 72 hours after infection. Una virus can be preserved by means of lyophilization of a brain suspension with and without fillers in a vacuum dryer. Of those we studied 5% peptone was the best filler.[stabilizer]

The heightened recent interest in investigation of arbor viruses is related to their isolation in many of the nations of the world [1, 5]. We made a study of the biological distinctions and developed optimum methods of passing and preserving Una arbor virus which is referable to group A arboviruses according to the Casals classification [6]. There is no detailed description of the properties of this virus in the literature.

the cellular layer. After maintaining the culture at 33° for four days these findings do not change. On the VNK-21 culture the CPD<sub>50</sub> constitutes 10<sup>-6,6</sup> - 10<sup>-6,6</sup>. The PP dosage on the CE culture constitutes 5·10 PPU/ml [plaque producing units per milliliter].

Table 1  
CPE and PP of Una virus in different tissue cultures

| a)<br>Культура ткани | b)<br>Цитопатический эффект | в) Ближкообразующая способность вируса |                    |
|----------------------|-----------------------------|--|--------------------|
|                      |                             | д) метод Портерфилда                   | е) метод Карповича |
| VNK-21               | +                           | - <sup>1</sup>                         | 0                  |
| CE                   | —                           | +                                      | +                  |
| HEK                  | —                           | 0                                      | 0                  |
| HEp-2                | —                           | 0                                      | 0                  |
| GMH                  | —                           | 0                                      | 0                  |

Legend:

- a) tissue culture
- b) CPE
- c) PP capacity of the virus
- d) Porterfield method
- e) Karpovich method

<sup>1</sup>The VNK-21 tissue culture is not suitable for demonstrating the plaque producing capacity of the virus since the cells degenerate within 24 hours under an agar cover.

Una virus was able to form plaques only on the CE culture. They were visible on the second postinfection day. Their number and appearance did not change upon further cultivation at 33°. The plaques are round, 1-1.5 mm in diameter, with irregular borders. The appearance of the plaques formed on CE is the same when using different covers, but they are less transparent when covered by the Karpovich method.

Table 2 shows that the virus appeared in the blood of mice only six hours after inoculation. Una virus is characterized by two rises in blood titers: six (10<sup>-2,2</sup>) and 48 (10<sup>-2,3</sup>) hours after infection. The virus was demonstrable in brain tissue three hours after inoculation and the quantity increased gradually. The highest titer was observed after 72 hours (10<sup>-4,3</sup>), thereafter it declined gradually. On the seventh day the titer was 10<sup>-1</sup>.

In observing the sick mice we noticed that they were much less active than the healthy ones, their fur was often ruffled, and paralysis of the hind extremities was observed on the 4th-5th postinoculation day. Most of the animals died on the 4th day after infection.

**Table 2**  
**Dynamics of accumulation of Una virus in the blood and brain**  
**tissue of infected mice**

| a) Источник вируса  | b) Сроки после заражения (в часах) |            |            |          |            |            |            |            |          |
|---------------------|------------------------------------|------------|------------|----------|------------|------------|------------|------------|----------|
|                     | 3                                  | 6          | 12         | 24       | 48         | 72         | 96         | 144        | 168      |
| Мозг c)<br>Кровь d) | 0,8*<br>0                          | 1,6<br>2,2 | 1,7<br>1,5 | 2,5<br>2 | 2,2<br>2,3 | 4,3<br>0,9 | 3,7<br>1,2 | 2,2<br>0,5 | 1<br>0,5 |

**Legend:**

- a) source of virus                      c) brain  
b) time after infection (hours)      d) blood

Note: the numbers in the table indicate the lg of viral titers

**Table 3**  
**Changes in viral titers after desiccation and storage of lyophilized suspensions at different temperatures**

| a   | Способ консервации | b До сушки | c После сушки | d Срок хранения |                   |     |          |     |           |     |     |     |  |  |  |
|---|--------------------|------------|---------------|-----------------|-------------------|-----|----------|-----|-----------|-----|-----|-----|--|--|--|
|   |                    |            |               | e 1 месяц       | 2 месяца          |     | 3 месяца |     | 7 месяцев |     |     |     |  |  |  |
|   |                    |            |               |                 |                   |     |          |     |           |     |     |     |  |  |  |
|   |                    |            |               |                 | f способ хранения |     |          |     |           |     |     |     |  |  |  |
| T   | K                  | Ф          | Z             | T               | K                 | T   | K        | Ф   | Z         | T   | K   |     |  |  |  |
| g Вирус на физиологическом растворе без наполнителя | 4,8                | 5,1        | 1,7           | 1               | 3,5               | 3   | 1        | 1   | 1         | 1,7 | 2,6 | 3   |  |  |  |
| h " + 10% сахара + 1% желатина                      | 4,3                | 4,7        | 2,8           | 3,0             | 2,5               | 3,7 | 1        | 3   | 1         | 2,7 | 3   | 4,1 |  |  |  |
| i " + 20% молока                                    | 4,6                | 5,4        | 4             | 6               | 4,5               | 4,3 | 3,8      | 4,3 | 1         | 3,5 | 4,4 | 3,1 |  |  |  |
| j " + 5% пептона                                    | 4,2                | 5          | 3,5           | 5,5             | 4,3               | 4,0 | 2        | 4,3 | 2         | 4,8 | 4,1 | 4,3 |  |  |  |

**Legend:**

- a) means of preservation      f) storage method  
b) before desiccation          g) virus on saline without stabilizer  
c) after "                      h) " + 10% saccharose + 1% gelatin  
d) storage time                i) " + 20% milk  
e) ... months                 j) " + 5% peptone

Note: the numbers in the table indicate the lg of viral titers



Table 3 shows the changes in viral titers in the course of drying and storage of lyophilized suspensions at different temperatures. As we see the virus containing suspension both with and without stabilizers tolerates the drying process well under the conditions chosen. A marked drop in titers was noted when Una virus, dried without stabilizers, was stored at thermal and room temperature for one month. The optimum stabilizer is peptone in an end concentration of 5%. It as well as 20% milk retarded the drop in viral titer while stored at 4° for the duration of the observation period. Virus stored at -20° without stabilizer and with stabilizers also retained its titers well during the observation period.

#### BIBLIOGRAPHY

1. V.M. Zhdanov, S.Ya. Gaydamovich, Virusologiya (Virology), Moscow, 1966.
2. L.G. Karpovich, in the book: Kleshchevov Entsefalit, Kemerovskaya Kleshchevaya Likhoradka, Gemorragicheskiye Likhoradki i Drugiye Arbovirusnyye Infektsii (Tick-borne Encephalitis, Kemerov Tick Fever, Hemorrhagic Fever and Other Arbovirus Infections), Moscow, 1964, p 44.
3. O.N. Shcheglovitova, in the book: Arbovirusy (Arboviruses), Moscow, 1967, p 160.
4. L.S. Yakovleva, in the book: Voprosy Med.Virusologii (Problems in Medical Virology), Moscow, No 6, 1959, p 264.
5. J. Casals, J.Exp.Med., Vol 99, 1954, p 429.
6. O.R. Causey, J. Casals, R. Shope et al., Am.J.Trop.Med.Hyg., Vol 12, 1963, p 777.
7. J.S. Porterfield, Nature, Vol 183, 1959, p 1069.

Submitted to press 29 December 1967

UNCLASSIFIED

Security Classification

## DOCUMENT CONTROL DATA - P &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

|  |                             |   |  |
|--|-----------------------------|---|--|
| 1. ORIGINATING ACTIVITY (Corporate author)<br>Foreign Science and Technology Center<br>US Army Materiel Command<br>Department of the Army  |                             | 22. REPORT SECURITY CLASSIFICATION<br><b>UNCLASSIFIED</b>                         |  |
| 3. REPORT TITLE<br><br>BIOLOGICAL CHARACTERISTICS OF UNA VIRUS   |                             | 23. GROUP   |  |
| 4. DESCRIPTIVE NOTES (Type of report and inclusive dates)<br>• Translation   |                             |   |  |
| 5. AUTHOR(S) (First name, middle initial, last name)<br><br>O. N. Shcheglovitova   |                             |   |  |
| 6. REPORT DATE<br>27 October 69  | 7A. TOTAL NO. OF PAGES<br>5 | 7B. NO. OF REFS<br>N/A  |  |
| 8A. CONTRACT OR GRANT NO.  |                             | 8B. ORIGINATOR'S REPORT NUMBER(S)   |  |
| A. PROJECT NO.   |                             | FSTC-HT-23-061-70   |  |
| C. 02RD500 2301  |                             | 9. OTHER REPORT NUM (Any other numbers that may be assigned this report)          |  |
| D. SMJED-PR-I  |                             | ACSI Control Number J-7191  |  |
| 10. DISTRIBUTION STATEMENT<br><br>This document has been approved for public release and sale; its distribution is unlimited.  |                             |   |  |
| 11. SUPPLEMENTARY NOTES  |                             | 12. SPONSORING MILITARY ACTIVITY<br>US Army Foreign Science and Technology Center |  |
| 13. ABSTRACT<br><br><p>Interest in arbor viruses has heightened recently in view of the isolation of such viruses in <sup>many</sup> of the world's countries. The present investigation deals with the biological distinctions and development of optimum methods for passing and preserving UNA arbor virus which is referable to group A according to the Casals classification. In the course of this study it was found that UNA virus induces CPE in VNK-21 tissue cultures and plaque production in chick fibroblast cultures. The clinical symptoms of pathology following infection of albino mice weighing 6-7 grams with UNA virus, intracerebrally, are characterized by acute development of disease, paralysis of the hind legs on the 4th-5th postinfection day. The virus is demonstrable in the blood in the highest titers 6 and 48 hours after infection, and in the brain 72 hours after infection. UNA virus can be preserved by means of lyophilization of a brain suspension with and without fillers in a vacuum dryer. Of those we studied 5% peptone was the best filler (stabilizer).</p> |                             |   |  |

DD FORM 1473

USE PREVIOUS EDITIONS FOR ARMY USE.

UNCLASSIFIED

**UNCLASSIFIED**  
Security Classification

| 14. KEY WORDS  | LINK A |    | LINK B |    | LINK C |    |
|--|--------|----|--------|----|--------|----|
|  | ROLE   | WT | ROLE   | WT | ROLE   | WT |
| UNA virus<br>CPE<br>VNK-21 tissue cultures<br>Plaque production<br>Fibroblast cultures<br>Lyophilization |        |    |        |    |        |    |

**UNCLASSIFIED**  
Security Classification